MICHIGAN DEPARTMENT OF ENVIRONMENTAL QUALITY

INTEROFFICE COMMUNICATION

November 10, 1998

TO: File for sodium iodide (CAS# 7681-82-5)

FROM: Cathy Simon, Supervisor, Toxics Unit, Air Quality Division

SUBJECT: Change in the Initial Threshold Screening Level (ITSL)

The ITSL for sodium iodide has been changed from 0.04 $\mu g/m^3$ to 0.1 $\mu g/m^3$ based on annual averaging time.

The change in the ITSL was made due to a revision in the State's air toxic rules which became effective on November 10, 1998. Previously, the ITSL had been set pursuant to Rule 232 (i). This rule sets the ITSL at a default value of $0.04 \, \mu g/m^3$ (annual average) when no specific data are available to determine an ITSL. The November 10, 1998 revisions to the rules changed this default ITSL to a value of $0.1 \, \mu g/m^3$.

No updated review of the literature has been done since the ITSL was originally set at a value of $0.04~\mu g/m^3$, to determine if new data are available for this compound.

CAS: SLB

MICHIGAN DEPARTMENT OF NATURAL RESOURCES

INTEROFFICE COMMUNICATION

November 1, 1993

TO: File for Sodium Iodide (CAS No. 7681-82-5)

FROM: Marco Bianchi

SUBJECT: Initial Threshold Screening Level

The initial threshold screening level (ITSL) for sodium iodide is $0.04 \ \mu g/m^3$ based on an annual averaging time.

The following references or databases were searched to identify data to determine the ITSL: IRIS, HEAST, NTP Management Status Report, RTECS, EPB-CCD, EPB library, CAS-online, NLM-online, IARC, NIOSH Pocket Guide, and ACGIH Guide.

A review of RTECs provided nonessential information about sodium iodide, and data obtained through CAS- and NLM-online was insufficient to derive an ITSL. No other information about this chemical was available, therefore, the ITSL is set at trace.

ITSL for sodium iodide = $0.04 \mu g/m^3$ based on annual averaging.

MB:ma