

**MICHIGAN DEPARTMENT OF ENVIRONMENTAL QUALITY**

---

**INTEROFFICE COMMUNICATION**

---

September 6, 2000

TO: File for Sodium Glucoheptonate (CAS # 31138-65-5)  
FROM: Michael Depa, Toxics Unit, Air Quality Division  
SUBJECT: Screening Level Determination

The initial threshold screening level (ITSL) for sodium glucoheptonate is  $0.1\mu\text{g}/\text{m}^3$  based on an annual averaging time.

The following references or databases were searched to identify data to determine the screening level: U.S. EPA Integrated Risk Information System (IRIS), Registry for Toxic Effects of Chemical Substances (RTECS), American Conference of Governmental and Industrial Hygienists (ACGIH) Threshold Limit Values (TLVs), National Institute for Occupational Safety and Health (NIOSH) Pocket Guide to Hazardous Chemicals, Environmental Protection Bureau Library, International Agency for Research on Cancer (IARC) Monographs, Chemical Abstract Service (CAS) Online (1967-April, 2000), National Library of Medicine (NLM), Health Effects Assessment Summary Tables (HEAST), and National Toxicology Program (NTP) Status Report. The EPA has not established a reference dose (RfD) or a reference concentration (RfC) for sodium glucoheptonate. There are no occupational exposure limits (e.g., ACGIH TLV or NIOSH REL). The molecular formula is  $\text{C}_7\text{-H}_{14}\text{-O}_8\text{.Na}$  with a molecular weight of 248g.

Since there was no adequate toxicological data with which to derive a screening level, Rule 232(1)i was used to establish the screening level of  $0.1\mu\text{g}/\text{m}^3$  with an annual averaging time.

MD:LH